

8oz front



ATHENA
MILDEW CONTROL
LIQUID CONCENTRATE

SUSTAINABLE
NON-BURNING
USE ANYTIME

Pure Mildew Control

Homeopathically Controls Downy and Powdery Mildew

Mildew Control is a fungicidal agent naturally derived from plant extracts. It is applied as a foliar spray and is effective against downy and powdery mildew.

Makes 6 gallons

ACTIVE INGREDIENT:
Geraniol 2.14%

INERT INGREDIENTS:* 97.86%

TOTAL: 100.00%

*Water, Isopropyl alcohol

CAUTION **KEEP OUT OF REACH OF CHILDREN**
See Back Panel for Precautionary Statements

NET CONTENTS: 8 fl oz (236 ml)

16oz front



ATHENA
MILDEW CONTROL
LIQUID CONCENTRATE

SUSTAINABLE
NON-BURNING
USE ANYTIME

*Pure
Mildew Control*

**Homeopathically Controls
Downy and Powdery Mildew**

Mildew Control is a fungicidal agent naturally derived from plant extracts. It is applied as a foliar spray and is effective against downy and powdery mildew.

Makes 12 gallons

ACTIVE INGREDIENT:
Geraniol 2.14%
INERT INGREDIENTS:* 97.86%
TOTAL: 100.00%
*Water, Isopropyl alcohol

CAUTION **KEEP OUT OF REACH OF CHILDREN**
See Back Panel for Precautionary Statements

NET CONTENTS: 16 fl oz (473 ml)

32oz front

 ATHENA
**MILDEW
CONTROL**

LIQUID CONCENTRATE

SUSTAINABLE
NON-BURNING
USE ANYTIME

*Pure
Mildew Control*

**Homeopathically Controls
Downy and Powdery Mildew**

Mildew Control is a fungicidal agent naturally derived from plant extracts. It is applied as a foliar spray and is effective against downy and powdery mildew.

Makes 25 gallons

ACTIVE INGREDIENT:

Geraniol 2.14%

INERT INGREDIENTS:* 97.86%

TOTAL: 100.00%

*Water, Isopropyl alcohol

CAUTION KEEP OUT OF REACH OF CHILDREN
See Back Panel for Precautionary Statements

NET CONTENTS: 32 fl oz (946 ml)

64oz front



ATHENA MILDEW CONTROL

LIQUID CONCENTRATE

*Pure
Mildew Control*

**Homeopathically Controls
Downy and Powdery Mildew**

Mildew Control is a fungicidal agent naturally derived from plant extracts. It is applied as a foliar spray and is effective against downy and powdery mildew.

SUSTAINABLE

NON-BURNING

USE ANYTIME

Makes 50 gallons

ACTIVE INGREDIENT:

Geraniol 2.14%

INERT INGREDIENTS:* 97.86%

TOTAL: 100.00%

*Water, Isopropyl alcohol

CAUTION KEEP OUT OF REACH OF CHILDREN
See Back Panel for Precautionary Statements

NET CONTENTS: 64 fl oz (1.89 L)

8oz back



MILDEW CONTROL

Pure Mildew Control

- Effective against *Podosphaera fusca*, the powdery mildew found on cucumbers, luffas, melons (including watermelons), and squashes (including pumpkins).
- Effective against *Erysiphe necator* (*Uncinula necator*), the powdery mildew endemic to grapes.
- Effective against *Podosphaera leucotricha*, the powdery mildew that affects both apples and pears.
- Effective against *Podosphaera aphanis*, the powdery mildew commonly found on rosaceae, such as blackberries, raspberries, and roses.
- Effective against *Oidium lycopersicum*, the powdery mildew of tomatoes.
- Effective against powdery mildew on ornamental and lawn and turf.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage and disposal. Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, and feed.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after

www.athenaproducts.com

Athena Products, Inc
8190 Mission Blvd, Ste B#7
Riverside, CA 92509-2999
Call toll free 1-844-333-1818



puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

PESTICIDES DISPOSAL: Reuse of this container is prohibited. Please dispose of this recyclable container at a container collection site, which can be found by contacting your municipality for the location nearest you.

ENVIRONMENTAL HAZARDS: For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Mix Rate: The mix rate for Mildew Control concentrate is 1.28 oz [37.9 ml] per one gallon of water. SHAKE OR MIX THOROUGHLY! Reverse Osmosis water is recommended.

Avoid using during peak sunlight or intense lighting, as some plants are sensitive to oil based products. The buyer undertakes all accountability for correct usage of this product.

Crop Name	Mix Rates	Dilution Rate	Foliar Spray	Timing & Frequency
All crops, except Strawberry*	1.28 fl oz [37.9 ml] / 1 gal	1:100	1.5 fl oz (45 ml) / 1,000R ² 70.3 fl oz (2,088 L) / acre 1.32 gal (5L) / ha	Low Severity disease outbreak
	1 gal [3.79 L] / 100 gal		2.4 fl oz (70 ml) / 1,000R ² 105.6 fl oz (3,131 L) / acre 1.98 gal (75L) / ha	High Severity disease outbreak

*Not recommended for use on strawberries, as it may cause phytotoxicity to flowers.

FIRST AID

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. **IF INHALED** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. **IF IN EYES** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment. **HOTLINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information call the National Pesticides Information Center at 1-800-858-7378, 6:30 AM to 4:30 PM Pacific time (PT), seven days a week. During other times, call the poison control center 1-800-222-1222.

This product has not been registered by the United States Environmental Protection Agency. Athena Products represents that this product qualifies for exemption from registration under FIFRA 25(b).

ESL25804A REV052018

16oz back



MILDEW CONTROL

Pure Mildew Control

- Effective against *Podosphaera fusca*, the powdery mildew found on cucumbers, luffas, melons (including watermelons), and squashes (including pumpkins).
- Effective against *Erysiphe necator (Uncinula necator)*, the powdery mildew endemic to grapes.
- Effective against *Podosphaera leucotricha*, the powdery mildew that affects both apples and pears.
- Effective against *Podosphaera aphanis*, the powdery mildew commonly found on rosaceae, such as blackberries, raspberries, and roses.
- Effective against *Didium lycopersicum*, the powdery mildew of tomatoes.
- Effective against powdery mildew on ornamental and lawn and turf.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage and disposal. Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, and feed.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

www.athenaproducts.com

Athena Products Inc
8190 Mission Blvd, Ste B#7
Riverside, CA 92509-2999
Call toll free 1-844-333-1818



6 94157 85865 4

PESTICIDES DISPOSAL: Reuse of this container is prohibited. Please dispose of this recyclable container at a container collection site, which can be found by contacting your municipality for the location nearest you.

ENVIRONMENTAL HAZARDS: For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

ESL25b04A REV052118

Mix Rate: The mix rate for Mildew Control concentrate is 1.28 oz (37.9 ml) per one gallon of water. SHAKE OR MIX THOROUGHLY! Reverse Osmosis water is recommended.

Avoid using during peak sunlight or intense lighting, as some plants are sensitive to oil based products. The buyer undertakes all accountability for correct usage of this product.

Crop Name	Mix Rates	Dilution Rate	Foliar Spray	Timing & Frequency
All crops, except Strawberry*	1.28 fl oz (37.9 ml) /1 gal	1:100	1.5 fl oz (45 ml)/1,000ft ² 70.3 fl oz (2.08L)/Acre 1.32 gal (5L)/Ha	Low Severity disease outbreak
	1 gal (3.79L) /100 gal		2.4 fl oz (70 ml)/1,000ft ² 105.6 fl oz (3.13L)/Acre 1.98 gal (75L)/Ha	High Severity disease outbreak

*Not recommended for use on strawberries, as it may cause phytotoxicity to flowers.

FIRST AID

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. **IF INHALED** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. **IF IN EYES** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment. **HOT LINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information call the National Pesticides Information Center at 1-800-858-7378, 8:30 AM to 4:30 PM Pacific time (PT), seven days a week. During other times, call the poison control center 1-800-222-1222.

This product has not been registered by the United States Environmental Protection Agency. Athena Products Inc represents that this product qualifies for exemption from registration under FIFRA 25(b).

32oz back



MILDEW CONTROL

Pure Mildew Control

- Effective against *Podosphaera fusca*, the powdery mildew found on cucumbers, luffas, melons (including watermelons), and squashes (including pumpkins).
- Effective against *Erysiphe necator (Uncinula necator)*, the powdery mildew endemic to grapes.
- Effective against *Podosphaera leucotricha*, the powdery mildew that affects both apples and pears.
- Effective against *Podosphaera aphanis*, the powdery mildew commonly found on rosaceae, such as blackberries, raspberries, and roses.
- Effective against *Oidium lycopersicum*, the powdery mildew of tomatoes.
- Effective against powdery mildew on ornamental and lawn and turf.

Mix Rate: The mix rate for Mildew Control concentrate is 1.28 fl oz (37.9 ml) per one gallon of water. SHAKE OR MIX THOROUGHLY! Reverse Osmosis water is recommended.

Avoid using during peak sunlight or intense lighting, as some plants are sensitive to oil based products. The buyer undertakes all accountability for correct usage of this product.

Crop Name	Mix Rates	Dilution Rate	Foliar Spray	Timing & Frequency
All crops, except Strawberry*	1.28 fl oz (37.9 ml) /1 gal	1:100	1.5 fl oz (45 ml)/1,000ft ² 70.3 fl oz (2.08L)/Acre 1.32 gal (5L)/Ha	Low Severity disease outbreak
	1 gal (3.79 L) /100 gal		2.4 fl oz (70 ml)/1,000ft ² 105.6 fl oz (3.13L)/Acre 1.98 gal (7.5L)/Ha	High Severity disease outbreak

*Not recommended for use on strawberries, as it may cause phytotoxicity to flowers.

FIRST AID

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. **IF INHALED** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. **IF IN EYES** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment. **HOTLINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information call the National Pesticides Information Center at 1-800-858-7378, 6:30 AM to 4:30 PM Pacific time (PT), seven days a week. During other times, call the poison control center 1-800-222-1222.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage and disposal. Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food and feed.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

PESTICIDES DISPOSAL: Reuse of this container is prohibited. Please dispose of this recyclable container at a container collection site, which can be found by contacting your municipality for the location nearest you.

ENVIRONMENTAL HAZARDS: For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

This product has not been registered by the United States Environmental Protection Agency. Athena Products Inc represents that this product qualifies for exemption from registration under FIFRA 25(b).

ESL25b04A REV052118

Athena Products Inc
8190 Mission Blvd, Ste B#7
Riverside, CA 92509-2999
Call toll free 1-844-333-1818

www.athenaproducts.com



6 94 157 85864 7

64oz back



MILDEW CONTROL

Pure Mildew Control

- Effective against *Podospaera fusca*, the powdery mildew found on cucumbers, luffas, melons (including watermelons), and squashes (including pumpkins).
- Effective against *Erysiphe necator (Uncinula necator)*, the powdery mildew endemic to grapes.
- Effective against *Podospaera leucotricha*, the powdery mildew that affects both apples and pears.
- Effective against *Podospaera aphanis*, the powdery mildew commonly found on rosaceae, such as blackberries, raspberries, and roses.
- Effective against *Didium lycopersicum*, the powdery mildew of tomatoes.
- Effective against powdery mildew on ornamental and lawn and turf.

Mix Rate: The mix rate for Mildew Control concentrate is 1.28 fl oz [37.9] ml per one gallon of water. SHAKE OR MIX THOROUGHLY! Reverse Osmosis water is recommended.

Avoid using during peak sunlight or intense lighting, as some plants are sensitive to oil based products. The buyer undertakes all accountability for correct usage of this product.

Crop Name	Mix Rates	Dilution Rate	Foliar Spray	Timing & Frequency
All crops, except Strawberry*	1.28 fl oz [37.9 ml] /1 gal	1:100	1.5 fl oz [45 ml]/1,000ft ² 70.3 fl oz [2.08L]/Acre 1.32 gal [5L]/Ha	Low Severity disease outbreak
	1 gal [3.79 L] /100 gal		2.4 fl oz [70 ml]/1,000ft ² 105.6 fl oz [3.13L]/Acre 1.98 gal [7.5L]/Ha	High Severity disease outbreak

*Not recommended for use on strawberries, as it may cause phytotoxicity to flowers.

FIRST AID

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. **IF INHALED** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. **IF IN EYES** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment. **HOT LINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information call the National Pesticides Information Center at 1-800-858-7378, 6:30 AM to 4:30 PM Pacific time (PT), seven days a week. During other times, call the poison control center 1-800-222-1222.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage and disposal. Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food and feed.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

PESTICIDES DISPOSAL: Reuse of this container is prohibited. Please dispose of this recyclable container at a container collection site, which can be found by contacting your municipality for the location nearest you.

ENVIRONMENTAL HAZARDS: For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

This product has not been registered by the United States Environmental Protection Agency. Athena Products Inc represents that this product qualifies for exemption from registration under FIFRA 25(b).

ESL25b04A REV052118

Athena Products Inc
8190 Mission Blvd, Ste B#7
Riverside, CA 92509-2999
Call toll free 1-844-333-1818

www.athenaproducts.com

